

On the Spectrum of Middle-Cubes*

Ke Qiu^a, Rong Qiu^b, Yong Jiang^b, Jian Shen^c

^aDepartment of Computer Science, Brock University, St. Catharines, Ontario, Canada

^bState Key Laboratory of Fire Science, University of Science and Technology of China

Hefei, Anhui 230026, P.R. China

^cDepartment of Mathematics, Texas State University, San Marcos, TX 78666, USA

Abstract

A middle-cube is an induced subgraph consisting of nodes at the middle two layers of a hypercube. The middle-cubes are related to the well-known Revolving Door (Middle Levels) conjecture. We study the middle-cube graph by completely characterizing its spectrum. Specifically, we first present a simple proof of its spectrum utilizing the fact that the graph is related to Johnson graphs which are distance-regular graphs and whose eigenvalues can be computed using the association schemes. We then give a second proof from a pure graph theory point of view without using its distance regular property and the technique of association schemes.

1 Introduction

The n -dimensional *hypercube*, Q_n , has 2^n nodes such that two nodes u and v , $0 \leq u, v \leq 2^n - 1$, are connected if and only if their binary representations differ in exactly one bit. For an odd $n = 2k + 1$, the *middle-cube*, M_n , is the subgraph induced by all the nodes whose binary representations have either k 1's or $k + 1$ 1's. Fig. 1 shows a 3-cube with its middle-cube highlighted.

The middle-cube M_{2k+1} consists of the nodes at the middle two layers of the corresponding hypercube Q_{2k+1} . Equivalently, these nodes are at the middle levels k and $k + 1$ of the Boolean lattice $\mathcal{B}_{2k+1}$ (and the Hasse diagram of $\mathcal{B}_{2k+1}$ is isomorphic to Q_{2k+1}) [10]. Middle-cubes have been considered as a possible topology to interconnect processors in networks [8]. A well-known open problem concerning middle-cubes is *the Revolving Door (Middle Levels) conjecture* [6, 14]: All middle-cubes M_{2k+1} are Hamiltonian. The conjecture has been verified for $k \leq 17$ [10] but remains open in general. Partial results on the conjecture can be found in [7, 9, 10]. In particular, Johnson proved in 2004 that M_{2k+1} has a cycle of length $(1 - o(1))|M_{2k+1}|$, where $|M_{2k+1}| = 2^{\binom{2k+1}{k}}$ is the number of vertices in M_{2k+1} .

*Research supported by National Natural Science Foundation of China (No. 50676091) and Program for New Century Excellent Talents in University (NCET-06-0546).




Figure 1: A 3-cube

The *spectrum* of a graph consists of all distinct eigenvalues and their respective multiplicities of the adjacency matrix of the graph. It is worth mentioning that the spectral and structural properties of a graph are related [3, 13]. For example, van den Heuvel [12] proved some necessary spectral conditions for a graph to be Hamiltonian. To better understand various properties for the middle-cubes, it may be necessary to study the spectrum for middle-cubes. In the next section, we give a complete characterization for the spectrum of the middle-cubes by giving two different proofs, one from the distance-regular graph point of view, and the other from a pure graph theory point of view.

2 Spectrum of Middle-Cubes

We always assume $n = 2k + 1$ throughout the paper. Without confusion from the context, we abuse the notation M_n for both the middle cube and its adjacency matrix. The eigenvalues and their corresponding multiplicities for M_n , $n = 3, 5, 7, 9$, are given in Table 1. In this section, we will prove that

Theorem 1 *The characteristic polynomial of M_n is*

$$|\lambda \mathbf{I} - M_n| = \prod_{i=1}^{k+1} (\lambda \pm i)^{\binom{n}{k+1-i} - \binom{n}{k-i}}.$$

We also note that the sequence 1, 2, 1, 4, 5, 1, 6, 14, 14, ... appears in the *The On-Line Encyclopedia of Integer Sequences* as sequence A050166 [11, 5].

Table 1: Eigenvalues of Middle-Cubes M_n for $n = 3, 5, 7, 9$.

	-5	-4	-3	-2	-1	1	2	3	4	5
3				1	2	2	1			
5			1	4	5	5	4	1		
7		1	6	14	14	14	14	6	1	
9	1	8	27	48	42	42	48	27	8	1

We will first consider the middle cube by relating it to Johnson graphs. Let X be a finite set and e an positive integer. The *Johnson graph of the e -sets in X* has a vertex set $\binom{X}{e}$, the set of all e -subsets of X (subsets of cardinality e). Two vertices u and v are adjacent whenever $|u \cap v| = e - 1$ [1]. Since Johnson graphs are distance regular, the eigenvalues of Johnson graphs can be computed using association schemes described in [1, Chapter 2] and [2, Pages 69-72].

Proof. Let M_{2k+1}^2 be the adjacency matrix of the middle-cube. Let $J(n, m)$ be the adjacency matrix of the Johnson graph with vertex set $\binom{[n]}{m}$, where two m -subsets are adjacent when they have exactly $m - 1$ elements in common. Then by the definition of M_{2k+1}^2 and $J(n, m)$,

$$M_{2k+1}^2 = \begin{bmatrix} J(2k+1, k+1) & O \\ O & J(2k+1, k+1) \end{bmatrix} + (k+1)I.$$

By [2, Page 79], $J(n, m)$ has eigenvalues $(m-i)(n-m-i) - i$ with multiplicity $\binom{n}{i} - \binom{n}{i-1}$. (The proof of the spectrum of $J(n, m)$ requires techniques using the association schemes.) Then M_{2k+1}^2 has eigenvalues

$$(k+1-i)(k-i) - i + (k+1) = (k+1-i)^2$$

with multiplicity $2 \left(\binom{n}{i} - \binom{n}{i-1} \right)$. Since M_{2k+1}^2 is bipartite, it has symmetric positive/negative eigenvalues. Therefore M_{2k+1}^2 has eigenvalues $\pm(k+1-i)$ with multiplicity $\binom{n}{i} - \binom{n}{i-1}$. $\square$

We now study the spectrum of the middle cube from a graph theoretical point of view. Before we give a second proof for Theorem 1 using graph theory, several definitions are in order. We denote by S the set $\{1, 2, \dots, n\}$. We use $A + B$ to denote the union of the sets A and B . Similarly, $A - B$ denotes the set of elements that are in A but not in B , while $|A|$ represents the size of the set A . An r -set is a set of size r . We denote by $\binom{S}{k}$ the set of all k -subsets of S . For the convenience of our proofs, we also view $\binom{S}{k} + \binom{S}{k+1}$ as the vertex set of the middle-cube M_n . Thus the edge set of M_n is induced by the inclusion relation; that is, two distinct vertices A, B in $\binom{S}{k} + \binom{S}{k+1}$ are adjacent if and only if $A \subset B$ or $B \subset A$.

For each positive integer i , let $A_1^{(i)}, A_2^{(i)}, \dots, A_{\binom{n}{i}}^{(i)}$ be an ordering of all $\binom{n}{i}$ i -subsets of S . Let r be a fixed positive integer with $r \leq k$. Let $x_1, \dots, x_{\binom{n}{r}}$ be real variables. Define a weight function:

$$f(A_i^{(r)}) = x_i, \quad 1 \leq i \leq \binom{n}{r}$$

subject to the following $\binom{n}{r-1}$ constraints

$$\sum_{i \notin R} f(R + \{i\}) = 0 \text{ for each } R \in \binom{S}{r-1}. \quad (1)$$

For each $A \subseteq S$, we define

$$f(A) = \sum_{A_i^{(r)} \in A} f(A_i^{(r)}). \quad (2)$$

(Thus $f(A) = 0$ whenever $|A| \leq r - 1$.) For each i with $r \leq i \leq k$, we define

$$V_i^{(r)} = \left\{ \left(f \left(A_1^{(i)} \right), \dots, f \left(A_{\binom{n}{i}}^{(i)} \right) \right) \right\}$$

and

$$V_{i,i+1}^{(r)} = \left\{ \left(f \left(A_1^{(i)} \right), \dots, f \left(A_{\binom{n}{i}}^{(i)} \right), f \left(A_1^{(i+1)} \right), \dots, f \left(A_{\binom{n}{i+1}}^{(i+1)} \right) \right) \right\}.$$

Then both $(V_i^{(r)}, +)$ and $(V_{i,i+1}^{(r)}, +)$ are vector spaces on reals. We will show that $V_{k,k+1}^{(r)}$ is the eigenspace with dimension $\binom{n}{r} - \binom{n}{r-1}$ corresponding to the eigenvalue $k + 1 - r$ for the matrix M_n .

Lemma 1 *Let A be a subset of S . Suppose $i \notin A$ and $j \in A$. Then*

$$f(A + \{i\}) = f(A) + \sum_{R \in \binom{A}{r-1}} f(R + \{i\})$$

and

$$f(A - \{j\}) = f(A) - \sum_{R \in \binom{A - \{j\}}{r-1}} f(R + \{j\}).$$

Proof.

$$\begin{aligned} f(A + \{i\}) &= \sum_{R \in \binom{A + \{i\}}{r}} f(R) \\ &= \sum_{i \notin R \in \binom{A + \{i\}}{r}} f(R) + \sum_{i \in R \in \binom{A + \{i\}}{r}} f(R) \\ &= \sum_{R \in \binom{A}{r}} f(R) + \sum_{R - \{i\} \in \binom{A}{r-1}} f(R) \\ &= f(A) + \sum_{R \in \binom{A}{r-1}} f(R + \{i\}). \end{aligned}$$

Similarly,

$$\begin{aligned} f(A - \{j\}) &= \sum_{R \in \binom{A - \{j\}}{r}} f(R) \\ &= \sum_{R \in \binom{A}{r}} f(R) - \sum_{j \in R \in \binom{A}{r}} f(R) \\ &= f(A) - \sum_{R - \{j\} \in \binom{A - \{j\}}{r-1}} f(R) \\ &= f(A) - \sum_{R \in \binom{A - \{j\}}{r-1}} f(R + \{j\}). \end{aligned}$$

□

Lemma 2 *Let A be a subset of S . Then*

$$\sum_{i \notin A} \sum_{R \in \binom{A}{r-1}} f(R + \{i\}) = -r f(A).$$

Proof. The lemma is trivial if $|A| \leq r - 1$ (in which case $f(A) = 0$). Suppose now $|A| \geq r$. By (1) and Lemma 1,

$$\begin{aligned}
0 &= \sum_{R \in \binom{A}{r-1}} \sum_{i \notin R} f(R + \{i\}) \\
&= \sum_{R \in \binom{A}{r-1}} \left(\sum_{i \in A-R} f(R + \{i\}) + \sum_{i \notin A} f(R + \{i\}) \right) \\
&= \sum_{R \in \binom{A}{r-1}} \sum_{R \subset B \in \binom{A}{r}} f(B) + \sum_{R \in \binom{A}{r-1}} \sum_{i \notin A} f(R + \{i\}) \\
&= \sum_{B \in \binom{A}{r}} \sum_{R \in \binom{B}{r-1}} f(B) + \sum_{i \notin A} \sum_{R \in \binom{A}{r-1}} f(R + \{i\}) \\
&= r \sum_{B \in \binom{A}{r}} f(B) + \sum_{i \notin A} \sum_{R \in \binom{A}{r-1}} f(R + \{i\}) \\
&= r f(A) + \sum_{i \notin A} \sum_{R \in \binom{A}{r-1}} f(R + \{i\}),
\end{aligned}$$

from which Lemma 2 follows. $\square$

Lemma 3 *Let A be a subset of S . Then*

$$\sum_{i \in A} \sum_{R \in \binom{A-\{i\}}{r-1}} f(R + \{i\}) = r f(A).$$

Proof. The lemma is trivial if $|A| \leq r - 1$ (in which case $f(A) = 0$). Suppose now $|A| \geq r$. By (1), Lemmas 1 and 2,

$$\begin{aligned}
0 &= \sum_{R \in \binom{A}{r-1}} \sum_{i \notin R} f(R + \{i\}) \\
&= \sum_{R \in \binom{A}{r-1}} \left(\sum_{i \notin A} f(R + \{i\}) + \sum_{i \in A-R} f(R + \{i\}) \right) \\
&= \sum_{R \in \binom{A}{r-1}} \sum_{i \notin A} f(R + \{i\}) + \sum_{R \in \binom{A}{r-1}} \sum_{i \in A-R} f(R + \{i\}) \\
&= \sum_{i \notin A} \sum_{R \in \binom{A}{r-1}} f(R + \{i\}) + \sum_{i \in A} \sum_{R \in \binom{A-\{i\}}{r-1}} f(R + \{i\}) \\
&= -r f(A) + \sum_{i \in A} \sum_{R \in \binom{A-\{i\}}{r-1}} f(R + \{i\}),
\end{aligned}$$

from which Lemma 3 follows. $\square$

Recall that $V_{k,k+1}^{(r)} = \left\{ \left(f \left(A_1^{(k)} \right), \dots, f \left(A_n^{(k)} \right), f \left(A_1^{(k+1)} \right), \dots, f \left(A_{k+1}^{(k+1)} \right) \right) \right\}$ is a vector space on reals. Let E_λ be the eigenspace corresponding to the eigenvalue λ for the matrix M_n .

Lemma 4 *Let $n = 2k + 1$ and $1 \leq r \leq k$. Then $V_{k,k+1}^{(r)} \subseteq E_{k+1-r}$.*

Proof. For any vertex $A \in \binom{S}{k} + \binom{S}{k+1}$ in the middle-cube M_n , let $\Gamma(A)$ be the neighbor set of A . Then

$$\Gamma(A) = \begin{cases} \{A + \{i\} : i \notin A\} & \text{if } A \in \binom{S}{k}; \\ \{A - \{i\} : i \in A\} & \text{if } A \in \binom{S}{k+1}. \end{cases}$$

Thus to prove the lemma, it suffices to prove the following two identities:

$$\sum_{B \in \Gamma(A)} f(B) = \sum_{i \notin A} f(A + \{i\}) = (k + 1 - r) f(A) \text{ for each } A \in \binom{S}{k} \quad (3)$$

$$\sum_{B \in \Gamma(A)} f(B) = \sum_{i \in A} f(A - \{i\}) = (k+1-r)f(A) \text{ for each } A \in \binom{S}{k+1} \quad (4)$$

Proof of (3): By Lemmas 1 and 2,

$$\begin{aligned} \sum_{i \notin A} f(A + \{i\}) &= \sum_{i \notin A} \left(f(A) + \sum_{R \in \binom{A}{r-1}} f(R + \{i\}) \right) \\ &= \sum_{i \notin A} f(A) + \sum_{i \notin A} \sum_{R \in \binom{A}{r-1}} f(R + \{i\}) \\ &= (n - |A|)f(A) - rf(A) \\ &= (n - |A| - r)f(A) = (k+1-r)f(A). \end{aligned}$$

Proof of (4): By Lemmas 1 and 3,

$$\begin{aligned} \sum_{i \in A} f(A - \{i\}) &= \sum_{i \in A} \left(f(A) - \sum_{R \in \binom{A - \{i\}}{r-1}} f(R + \{i\}) \right) \\ &= \sum_{i \in A} f(A) - \sum_{i \in A} \sum_{R \in \binom{A - \{i\}}{r-1}} f(R + \{i\}) \\ &= |A|f(A) - rf(A) \\ &= (k+1-r)f(A). \end{aligned}$$

□

Lemma 5 *Let $n = 2k+1$ and $1 \leq r \leq k$. Then*

$$\dim V_{k,k+1}^{(r)} = \binom{n}{r} - \binom{n}{r-1}.$$

Proof. Let $M_{i,j}$ be the incidence matrix whose rows correspond to the i -subsets $A_1^{(i)}, A_2^{(i)}, \dots, A_{\binom{n}{i}}^{(i)}$, and whose columns correspond to the j -subsets $A_1^{(j)}, A_2^{(j)}, \dots, A_{\binom{n}{j}}^{(j)}$; that is, the (r, s) -entry of $M_{i,j}$ is 1 if $A_r^{(i)} \subset A_s^{(j)}$ or $A_s^{(j)} \subset A_r^{(i)}$, and 0 otherwise. By [4, Corollary 2], the matrix $M_{i,j}$ has full rank; that is,

$$\text{rank } M_{i,j} = \min \left\{ \binom{n}{i}, \binom{n}{j} \right\}.$$

Recall the definition that $V_r^{(r)} = \left\{ \left(f(A_1^{(r)}), \dots, f(A_{\binom{n}{r}}^{(r)}) \right) \right\}$. By (1), $V_r^{(r)}$ consists of all solution sets to the following homogeneous matrix equation:

$$(x_1, x_2, \dots, x_{\binom{n}{r}}) M_{r,r-1} = (0, 0, \dots, 0).$$

Thus

$$\dim V_r^{(r)} = \binom{n}{r} - \text{rank } M_{r,r-1} = \binom{n}{r} - \binom{n}{r-1}.$$

By (2) and the definition of $V_{k,k+1}^{(r)}$, each vector in $V_{k,k+1}^{(r)}$ can be written as

$$(x_1, x_2, \dots, x_{\binom{n}{r}}) \begin{bmatrix} M_{r,k} \\ \vdots \\ M_{r,k+1} \end{bmatrix}$$

for some vector $(x_1, x_2, \dots, x_{\binom{n}{r}}) \in V_r^{(r)}$. This implies that $V_{k,k+1}^{(r)} = V_r^{(r)} \begin{bmatrix} M_{r,k} & \vdots & M_{r,k+1} \end{bmatrix}$. Thus

$$\begin{aligned} \dim V_r^{(r)} &\geq \dim V_{k,k+1}^{(r)} \geq \dim V_r^{(r)} + \text{rank} \begin{bmatrix} M_{r,k} & \vdots & M_{r,k+1} \end{bmatrix} - \binom{n}{r} \\ &\geq \dim V_r^{(r)} + \text{rank } M_{r,k} - \binom{n}{r} \\ &= \dim V_r^{(r)} + \min \left\{ \binom{n}{r}, \binom{n}{k} \right\} - \binom{n}{r} \\ &= \dim V_r^{(r)} \end{aligned}$$

and so

$$\dim V_{k,k+1}^{(r)} = \dim V_r^{(r)} = \binom{n}{r} - \binom{n}{r-1}.$$

□

Theorem 2 *Let $n = 2k + 1$ and $1 \leq r \leq k$. Then*

$$E_{k+1-r} = V_{k,k+1}^{(r)}$$

and

$$\dim E_{k+1-r} = \dim E_{r-k-1} = \binom{n}{r} - \binom{n}{r-1}.$$

Furthermore, the characteristic polynomial of the matrix M_n is

$$|\lambda \mathbf{I} - M_n| = \prod_{i=1}^{k+1} (\lambda \pm i)^{\binom{n}{k+1-i} - \binom{n}{k-i}}.$$

Proof. The equation $\dim E_{k+1-r} = \dim E_{r-k-1}$ holds since the middle-cube M_n is a bipartite graph. Since M_n is a connected $(k+1)$ -regular graph, we have $\dim E_{k+1} = 1$. By Lemmas 4 and 5,

$$\begin{aligned} \binom{n}{k} + \binom{n}{k+1} &\geq \sum_{r=0}^k (\dim E_{k+1-r} + \dim E_{r-k-1}) \\ &= 2 \sum_{r=0}^k \dim E_{k+1-r} \\ &\geq 2 + 2 \sum_{r=1}^k \dim V_{k,k+1}^{(r)} \\ &= 2 + 2 \sum_{r=1}^k \left(\binom{n}{r} - \binom{n}{r-1} \right) \\ &= 2 \binom{n}{k} = \binom{n}{k} + \binom{n}{k+1}. \end{aligned}$$

Thus all equalities hold throughout. This also implies that all eigenvalues of M_n are integers i with $1 \leq |i| \leq k+1$, and that each eigenvalue of i has multiplicity $\binom{n}{k+1-i} - \binom{n}{k-i}$, where $\binom{n}{-1} = 0$. □

3 Conclusion

We prove that the characteristic polynomial of the middle-cube M_n with $n = 2k + 1$ is

$$\prod_{i=1}^{k+1} (\lambda \pm i)^{\binom{n}{k+1-i} - \binom{n}{k-i}}.$$

This spectral property may be useful in future research on various properties of the middle-cubes.

References

- [1] A. E. Brouwer, A. M. Cohen, and A. Neumaier, *Distance-Regular Graphs*, (Springer-Verlag, 1989).
- [2] A. E. Brouwer and W. H. Haemers, Spectra of Graphs, (*manuscript*, <http://www.cwi.nl/~aeb/math/ipm.pdf>).
- [3] D. M. Cvetković, M. Doob, and H. Sachs, *Spectra of Graphs*, 3rd ed. (Johann Ambrosius Barth, 1995).
- [4] D. H. Gottlieb, A certain class of incidence matrices, *Proc. Amer. Math. Soc.* **17** (1966), 1233–1237.
- [5] R. K. Guy, Catwalks, sandsteps and Pascal pyramids, *J. Integer Seq.* **3** (2000), no. 1, Article 00.1.6, (<http://www.cs.uwaterloo.ca/journals/JIS/VOL3/GUY/catwalks.html>).
- [6] I. Havel, Semipaths in directed cubes, in: M. Fiedler (Ed.), *Graphs and other Combinatorial Topics (Prague, 1982)*, 101–108, Teubner-Texte Math., 59, Teubner, Leipzig, 1983.
- [7] J. R. Johnson, Long cycles in the middle two layers of the discrete cube, *J. Combin. Theory Ser. A*, **105** (2004) 255–271.
- [8] S. V. R. Madabhushi, S. Lakshmivarahan, and S. K. Dhall, Analysis of the modified even networks, *Proc. of the 3rd IEEE Symposium on Parallel and Distributed Processing*, Dallas, Texas (1991) 128–131.
- [9] C. D. Savage and P. Winkler, Monotone Gray codes and the middle levels problem, *J. Combin. Theory Ser. A*, **70** (1995), 230–248.
- [10] I. Shields, B. J. Shields, and C. D. Savage, An update on the middle levels problem, *preprint* (2006).
- [11] N. J. A. Sloane, *The on-line encyclopedia of integer sequences*, (www.research.att.com/~njas/sequences/).
- [12] J. van den Heuvel, Hamilton cycles and eigenvalues of graphs, *Linear Algebra Appl.* **226/228** (1995), 723–730.
- [13] D. B. West, *Introduction to Graph Theory*, 2nd ed. (Prentice Hall, 2001).
- [14] D. B. West, Open problems - graph theory and combinatorics, (<http://www.math.uiuc.edu/~west/openp/revolving.html>).